arXiv:1703.09196v2 [math.CO] 28 Mar 2017

PATTERN RECOGNITION ON ORIENTED MATROIDS: DECOMPOSITIONS OF TOPES, AND ORTHOGONALITY RELATIONS

ANDREY O. MATVEEV

Abstract. If $\mathrm{V}(\boldsymbol{R})$ is the vertex set of a symmetric cycle $\boldsymbol{R}$ in the tope graph of a simple oriented matroid $\mathcal{M}$, then for any tope T of $\mathcal{M}$ there exists a unique inclusion-minimal subset $\boldsymbol{Q}(T,\boldsymbol{R})$ of $\mathrm{V}(\boldsymbol{R})$ such that T is the sum of the topes of $\boldsymbol{Q}(T,\boldsymbol{R})$.

If for decompositions $\boldsymbol{Q}(T',\boldsymbol{R}')$ and $\boldsymbol{Q}(T'',\boldsymbol{R}'')$ with respect to symmetric cycles $\boldsymbol{R}'$ and $\boldsymbol{R}''$ in the tope graphs of two simple oriented matroids, whose ground sets have the cardinalities of opposite parity, we have $|\boldsymbol{Q}(T',\boldsymbol{R}')|\geq 5$ and $|\boldsymbol{Q}(T'',\boldsymbol{R}'')|\geq 5$, then these decompositions satisfy a certain orthogonality relation.

1. Introduction

Let $\mathcal{M}':=(E_s,\mathcal{L}')=(E_s,\mathcal{T}')$ and $\mathcal{M}'':=(E_t,\mathcal{L}'')=(E_t,\mathcal{T}'')$ be *simple* (i.e., with no *loops*, *parallel elements* or *antiparallel elements*) oriented matroids on their ground sets $E_s=[s]:=\{1,\ldots,s\}$ and E_t, with sets of covectors $\mathcal{L}'$ and $\mathcal{L}''$, and with sets of topes $\mathcal{T}'\subseteq\{1,-1\}^{E_s}$ and $\mathcal{T}''\subseteq\{1,-1\}^{E_t}$, respectively; see [1] on oriented matroids. We suppose that

$$\begin{gathered}s<t\,,\\ s\not\equiv t\pmod 2\,.\end{gathered}\tag{1.1}$$

Let $\boldsymbol{R}'$ be a *symmetric* $2s$*-cycle*, with its vertex set $\mathrm{V}(\boldsymbol{R}')=-\mathrm{V}(\boldsymbol{R}')$, in the tope graph $\mathcal{T}'(\mathcal{L}')$ of $\mathcal{M}'$, and let $\boldsymbol{R}''$ be a *symmetric* $2t$*-cycle* in the tope graph $\mathcal{T}''(\mathcal{L}'')$ of $\mathcal{M}''$.

Let $T'\in\mathcal{T}'$ and $T''\in\mathcal{T}''$ be any topes of the oriented matroids $\mathcal{M}'$ and $\mathcal{M}''$ such that for the *unique inclusion-minimal subsets* $\boldsymbol{Q}(T',\boldsymbol{R}')\subset\mathrm{V}(\boldsymbol{R}')$ and $\boldsymbol{Q}(T'',\boldsymbol{R}'')\subset\mathrm{V}(\boldsymbol{R}'')$ with the properties

$$\sum_{Q'\in\boldsymbol{Q}(T',\boldsymbol{R}')}Q'=T'\quad\text{and}\quad\sum_{Q''\in\boldsymbol{Q}(T'',\boldsymbol{R}'')}Q''=T''\,,$$

discussed in [4, Sect. 3.11], we have

$$|\boldsymbol{Q}(T',\boldsymbol{R}')|\geq 5\quad\text{and}\quad|\boldsymbol{Q}(T'',\boldsymbol{R}'')|\geq 5\,.$$

Let

$$\boldsymbol{\Lambda}'':=\boldsymbol{\Lambda}(T'',\boldsymbol{R}'')$$

be the abstract simplicial complex on the vertex set E_t, with the facet family

$$\bigl\{E_t-\mathbf{S}(T'',Q'')\colon Q''\in\boldsymbol{Q}(T'',\boldsymbol{R}'')\bigr\}\,,\tag{1.2}$$

where $\mathbf{S}(T'', Q'')$ is the *separation set* of the topes T'' and Q''; see e.g. [5] on such complexes.

Associate with the complex $\boldsymbol{\Lambda}''$ its "*long*" *f-vector*

$$\boldsymbol{f}(\boldsymbol{\Lambda}''; t) := \big(f_0(\boldsymbol{\Lambda}''; t), f_1(\boldsymbol{\Lambda}''; t), \ldots, f_t(\boldsymbol{\Lambda}''; t)\big) \in \mathbb{N}^{t+1}$$

defined by

$$f_j(\boldsymbol{\Lambda}''; t) := \#\{F \in \boldsymbol{\Lambda}'' \colon |F| = j\}\ , \quad 0 \le j \le t\ ;$$

throughout the paper, the components of all vectors, as well as the rows and columns of matrices are indexed starting with zero.

Define vectors $\boldsymbol{\beta}(t; t) \in \mathbb{P}^{t+1}$ and $\boldsymbol{\beta}(s; t) \in \mathbb{N}^{t+1}$ by

$$\boldsymbol{\beta}(t; t) := \big(\tbinom{t}{0}, \tbinom{t}{1}, \ldots, \tbinom{t}{t}\big)\ ,$$
$$\boldsymbol{\beta}(s; t) := \big(\tbinom{s}{0}, \tbinom{s}{1}, \ldots, \tbinom{s}{t}\big)\ .$$

Let $\mathbf{U}(t)$ denote the square *backward identity matrix* of order $t+1$, whose (i, j)th entry is the Kronecker delta $\delta_{i+j,t}$. We denote by $\mathbf{T}(t)$ the square *forward shift matrix* of order $t+1$, whose (i, j)th entry is $\delta_{j-i,1}$.

According to the argument given in [5], and by [2, Prop. 3.51(a)], the vector

$$\big(\boldsymbol{\beta}(t; t) - \boldsymbol{f}(\boldsymbol{\Lambda}''; t)\big)\mathbf{U}(t) \in \mathbb{N}^{t+1}$$

is the *long f-vector* $\boldsymbol{f}(\boldsymbol{\Omega}''; t)$ of the *boundary complex* $\boldsymbol{\Omega}''$ of a $(t-3)$-dimensional *simplicial* convex *polytope* with t vertices.

Associate with the abstract simplicial complex

$$\boldsymbol{\Lambda}' := \boldsymbol{\Lambda}(T', \boldsymbol{R}')$$

on the vertex set E_s, with the facet family

$$\big\{E_s - \mathbf{S}(T', Q') \colon Q' \in \boldsymbol{Q}(T', \boldsymbol{R}')\big\}\ , \tag{1.3}$$

its *long f-vector*

$$\boldsymbol{f}(\boldsymbol{\Lambda}'; t) \in \mathbb{N}^{t+1}$$

defined by $f_j(\boldsymbol{\Lambda}'; t) := \#\{F \in \boldsymbol{\Lambda}' \colon |F| = j\}$, $0 \le j \le t$.

The vector

$$\big(\boldsymbol{\beta}(s; t) - \boldsymbol{f}(\boldsymbol{\Lambda}'; t)\big)\mathbf{T}(t)^{t-s}\mathbf{U}(t) \in \mathbb{N}^{t+1}$$

is the *long f-vector* $\boldsymbol{f}(\boldsymbol{\Omega}'; t)$ of the *boundary complex* $\boldsymbol{\Omega}'$ of an $(s-3)$-dimensional *simplicial* convex *polytope* with s vertices.

Define the (i, j)th entry of a square matrix

$$\mathbf{S}(t)$$

of order $t+1$ to be $(-1)^{j-i}\binom{t-i}{j-i}$.

Recall that the standard *h-vectors* $\boldsymbol{h}(\boldsymbol{\Omega}'') \in \mathbb{N}^{t-2}$ and $\boldsymbol{h}(\boldsymbol{\Omega}') \in \mathbb{N}^{s-2}$ of the *boundary complexes* $\boldsymbol{\Omega}''$ and $\boldsymbol{\Omega}'$ of *simplicial polytopes* both satisfy the *Dehn–Sommerville relations* [6, Sect. 8.3]

$$h_k(\boldsymbol{\Omega}'') = h_{t-k-3}(\boldsymbol{\Omega}'')\ , \qquad 0 \le k \le t-3\ ;$$
$$h_k(\boldsymbol{\Omega}') = h_{s-k-3}(\boldsymbol{\Omega}')\ , \qquad 0 \le k \le s-3\ .$$

As a consequence, the "*long*" *h-vectors*

$$\boldsymbol{h}(\boldsymbol{\Omega}'';t) := \big(\boldsymbol{\beta}(t;t) - \boldsymbol{f}(\boldsymbol{\Lambda}'';t)\big)\mathbf{U}(t)\mathbf{S}(t) \in \mathbb{Z}^{t+1} \tag{1.4}$$

and

$$\boldsymbol{h}(\boldsymbol{\Omega}';t) := \big(\boldsymbol{\beta}(s;t) - \boldsymbol{f}(\boldsymbol{\Lambda}';t)\big)\mathbf{T}(t)^{t-s}\mathbf{U}(t)\mathbf{S}(t) \in \mathbb{Z}^{t+1} \tag{1.5}$$

of the complexes $\boldsymbol{\Omega}''$ and $\boldsymbol{\Omega}'$ satisfy the *Dehn–Sommerville* type *relations*

$$h_k(\boldsymbol{\Omega}'';t) = -h_{t-k}(\boldsymbol{\Omega}'';t)\ , \qquad 0 \le k \le t\ ; \tag{1.6}$$

$$h_k(\boldsymbol{\Omega}';t) = \quad h_{t-k}(\boldsymbol{\Omega}';t)\ , \qquad 0 \le k \le t\ ; \tag{1.7}$$

see e.g. [4, Sect. 2.3].

2. Orthogonality relations for decompositions of topes with respect to symmetric cycles in the tope graphs

Since the maximal face $[t]$ of the simplex $\mathbf{2}^{[t]}$ does not belong to the complexes $\boldsymbol{\Omega}'$ and $\boldsymbol{\Omega}''$, by [4, Eq. (2.3) of Prop. 2.1] we have

$$\big\langle \boldsymbol{h}(\boldsymbol{\Omega}';t), \boldsymbol{\iota}(t)\big\rangle = \big\langle \boldsymbol{h}(\boldsymbol{\Omega}'';t), \boldsymbol{\iota}(t)\big\rangle = 0\ ,$$

for the vector

$$\boldsymbol{\iota}(t) := (1, 1, \ldots, 1) \in \mathbb{N}^{t+1}\ ,$$

where $\langle\cdot,\cdot\rangle$ means the standard scalar product.

For the positive integers k, define abstract simplicial complexes $\overline{\mathbf{2}^{[k]}}$ with their *long h-vectors* $\boldsymbol{h}(\overline{\mathbf{2}^{[k]}};t) := \boldsymbol{f}(\overline{\mathbf{2}^{[k]}};t)\mathbf{S}(t)$ to be the *boundary complexes* of the *simplices* $\mathbf{2}^{[k]}$ by

$$\overline{\mathbf{2}^{[k]}} := \mathbf{2}^{[k]} - \{[k]\}\ .$$

In view of (1.7), the long h-vector $\boldsymbol{h}(\boldsymbol{\Omega}')$ lies either in the linear span

$$\operatorname{span}\Big(\boldsymbol{h}(\overline{\mathbf{2}^{[1]}};t), \boldsymbol{h}(\overline{\mathbf{2}^{[3]}};t), \ldots, \boldsymbol{h}(\overline{\mathbf{2}^{[s-2]}};t)\Big)\ , \tag{2.1}$$

when t is *even*, or in the linear span

$$\operatorname{span}\Big(\boldsymbol{h}(\overline{\mathbf{2}^{[2]}};t), \boldsymbol{h}(\overline{\mathbf{2}^{[4]}};t), \ldots, \boldsymbol{h}(\overline{\mathbf{2}^{[s-2]}};t)\Big)\ , \tag{2.2}$$

when t is *odd*; see [4, Sect. 3.1].

The Dehn–Sommerville type relations (1.6) and (1.7) imply that $\boldsymbol{h}(\boldsymbol{\Omega}';t)$ is a *left eigenvector* of the backward identity matrix $\mathbf{U}(t)$ that corresponds to its eigenvalue 1, while $\boldsymbol{h}(\boldsymbol{\Omega}'';t)$ is a *right eigenvector* of $\mathbf{U}(t)$ that corresponds to the other eigenvalue -1. By the *principle of biorthogonality* [3, Th. 1.4.7(a)] we have

$$\big\langle \boldsymbol{h}(\boldsymbol{\Omega}';t), \boldsymbol{h}(\boldsymbol{\Omega}'';t)\big\rangle = 0\ . \tag{2.3}$$

In other words, together with the relation (2.3), the definitions (1.4) and (1.5) yield

$$\big(\boldsymbol{\beta}(s;t) - \boldsymbol{f}(\boldsymbol{\Lambda}';t)\big)\mathbf{T}(t)^{t-s}\mathbf{U}(t)\mathbf{S}(t)\mathbf{S}(t)^{\top}\mathbf{U}(t)\big(\boldsymbol{\beta}(t;t)^{\top} - \boldsymbol{f}(\boldsymbol{\Lambda}'';t)^{\top}\big) = 0\ .$$

Note that the (i,j)th entry of the square matrix

$$\mathbf{M}(t) := \mathbf{U}(t)\mathbf{S}(t)\mathbf{S}(t)^{\top}\mathbf{U}(t)$$

of order $t+1$ is $(-1)^{i+j}\binom{i+j}{i}$.

Let us sum up our conclusions:

Proposition 2.1. *Let $\mathcal{M}' := (E_s, \mathcal{L}') = (E_s, \mathcal{T}')$ and $\mathcal{M}'' := (E_t, \mathcal{L}'') = (E_t, \mathcal{T}'')$ be simple oriented matroids on their ground sets E_s and E_t such that*

$$s < t \,,$$
$$s \not\equiv t \pmod 2 \,,$$

with sets of covectors $\mathcal{L}'$ and $\mathcal{L}''$, and with sets of topes $\mathcal{T}' \subseteq \{1,-1\}^{E_s}$ and $\mathcal{T}'' \subseteq \{1,-1\}^{E_t}$, respectively.

Let $\boldsymbol{R}'$ be a symmetric cycle in the tope graph $\mathcal{T}'(\mathcal{L}')$ of $\mathcal{M}'$, and let $\boldsymbol{R}''$ be a symmetric cycle in the tope graph $\mathcal{T}''(\mathcal{L}'')$ of $\mathcal{M}''$.

Let $T' \in \mathcal{T}'$ and $T'' \in \mathcal{T}''$ be any topes of the oriented matroids $\mathcal{M}'$ and $\mathcal{M}''$ such that for the unique inclusion-minimal subsets $\boldsymbol{Q}(T', \boldsymbol{R}') \subset \mathrm{V}(\boldsymbol{R}')$ and $\boldsymbol{Q}(T'', \boldsymbol{R}'') \subset \mathrm{V}(\boldsymbol{R}'')$ with the properties

$$\sum_{Q' \in \boldsymbol{Q}(T', \boldsymbol{R}')} Q' = T' \quad \text{and} \quad \sum_{Q'' \in \boldsymbol{Q}(T'', \boldsymbol{R}'')} Q'' = T'' \,,$$

we have

$$|\boldsymbol{Q}(T', \boldsymbol{R}')| \geq 5 \quad \text{and} \quad |\boldsymbol{Q}(T'', \boldsymbol{R}'')| \geq 5 \,.$$

The long f-vectors $\boldsymbol{f}(\boldsymbol{\Lambda}''; t)$ and $\boldsymbol{f}(\boldsymbol{\Lambda}'; t)$ of the complexes $\boldsymbol{\Lambda}''$ and $\boldsymbol{\Lambda}'$ whose families of facets are defined by (1.2) *and* (1.3)*, respectively, satisfy the* orthogonality relation

$$\big(\boldsymbol{\beta}(s;t) - \boldsymbol{f}(\boldsymbol{\Lambda}'; t)\big)\mathbf{T}(t)^{t-s} \cdot \mathbf{M}(t) \cdot \big(\boldsymbol{\beta}(t;t)^{\top} - \boldsymbol{f}(\boldsymbol{\Lambda}''; t)^{\top}\big) = 0 \,.$$

References

[1] *Björner A., Las Vergnas M., Sturmfels B., White N., Ziegler G.M.* Oriented matroids. Second edition. Encyclopedia of Mathematics, 46. – Cambridge: Cambridge University Press, 1999.

[2] *Gainanov D.N.* Graphs for pattern recognition. Infeasible systems of linear inequalities. – Berlin: De Gruyter, 2016.

[3] *Horn R.A., Johnson C.R.* Matrix analysis. Second edition. – Cambridge: Cambridge University Press, 2013.

[4] *Matveev A.O.* Pattern recognition on oriented matroids. – Berlin: De Gruyter, 2017.

[5] *Matveev A.O.* Pattern recognition on oriented matroids: Decompositions of topes, and Dehn–Sommerville type relations. Preprint [arXiv:1703.04508], 2017.

[6] *Ziegler G.M.* Lectures on polytopes. Revised edition. Graduate Texts in Mathematics, 152. – Berlin: Springer-Verlag, 1998.

E-mail address: andrey.o.matveev@gmail.com